

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/28, C07K 1/00, C07H 21/02, 21/04, C12N 5/00, 15/00, G01N 33/53	A1	(11) International Publication Number: WO 97/16206 (43) International Publication Date: 9 May 1997 (09.05.97)
(21) International Application Number: PCT/US96/17119 (22) International Filing Date: 1 November 1996 (01.11.96) (30) Priority Data: 60/006,226 3 November 1995 (03.11.95) US (71)(72) Applicants and Inventors: GETZENBERG, Robert, H. [US/US]; 1627 Augusta Drive, Pittsburgh, PA 15237 (US). BAHNSON, Robert, R. [US/US]; 5200 Pembroke Place, Pittsburgh, PA 15232-1429 (US). (74) Agents: BENT, Stephen, A. et al.; Foley & Lardner, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US).	(81) Designated States: AU, CA, IL, JP, KR, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>	
(54) Title: BLADDER NUCLEAR MATRIX PROTEINS AND THEIR USE IN DETECTING AND TREATING CELL PROLIFERATIVE DISORDERS (57) Abstract Nuclear matrix proteins (NMPs) which are characterized by a defined expression in bladder tissue are provided. These NMPs are useful markers in diagnosing and monitoring the stage of malignancy, and treating cell proliferative disorders of the bladder. Also provided are substantially purified polypeptides and polypeptide sequences encoding the NMPs of the invention.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

WO 97/16206

PCT/US96/17119

**BLADDER NUCLEAR MATRIX PROTEINS AND THEIR USE IN DETECTING AND TREATING
CELL PROLIFERATIVE DISORDERS**

This invention was made with support from the University of Pittsburgh Cancer Institute and from NIH grant P30 CA47904 to the University of Pittsburgh Cancer Institute.

5

BACKGROUND OF THE INVENTION

The present invention relates generally to bladder nuclear matrix proteins, called "NMPs" here, and more specifically to novel nuclear matrix proteins of the bladder which are associated with cell-proliferative disorders.

The early diagnosis of bladder cancer is central to the effective treatment of the disease. Currently, there are no methods available to easily and specifically identify the presence of bladder cancer cells. The prevailing technique for diagnosis of bladder cancer is to identify bladder cancer cells by morphological examination of the cells by a pathologist. A cellular hallmark of the transformed phenotype is abnormal nuclear shape, the presence of multiple nucleoli and altered patterns of chromatin organization. Nuclear structural alterations are so prevalent in cancer cells that they are commonly used as a pathological marker of transformation for many types of cancer. Nuclear shape is determined in part by the nuclear matrix, the dynamic skeleton of the nucleus.

The nuclear matrix is the structural component of the nucleus that determines nuclear morphology, organizes the DNA in a three-dimensional fashion that is tissue specific, and has a central role in the regulation of a number of nuclear processes including the regulation of

WO 97/16206

-2-

PCT/US96/17119

gene expression. The nuclear matrix has been demonstrated to play a central role in the regulation of important cellular processes such as DNA replication and transcription. Getzenberg, *J. Cell Biochem.* 55: 22-31 (1994). The nuclear matrix is the framework or scaffolding of the nucleus and consists of the peripheral lamins and pore complexes, an internal ribonucleic protein network, and residual nucleoli. Berezney et al., *Biochem. Biophys. Res. Comm.* 60: 1410-17 (1974). The nuclear matrix consists of approximately 10% of the nuclear proteins and is virtually devoid of lipids, DNA and histones. Fey et al., *Critical Reviews in Eukaryotic Gene Expression* 1: 127-44 (1991).

A majority of the known NMPs are common to all cell types and physiologic states. A number of laboratories have identified NMPs which may be unique to certain cell types or states. Mitogenic stimulation and the induction of differentiation have been demonstrated to alter the composition of nuclear matrix proteins and structure. The nuclear matrix contains a number of associated proteins that have been demonstrated to be involved in transformation. Berezney first showed that the nuclear matrix is altered in transformation, examining hepatoma nuclear matrix proteins. Berezney et al., *Cancer Res.* 39: 3031-39 (1979). Fey and Penman demonstrated that tumor promoters induce a specific morphologic signature in the nuclear matrix-intermediate filament scaffold of kidney cells. Fey et al., *Proc. Nat'l Acad. Sci. USA* 81: 859-66 (1984). Fey and Penman went on to demonstrate that the pattern of NMPs differed between normal and tumorigenic cell lines. Fey et al., *loc. cit.* 85: 121-25 (1989). Recently, an antibody to a nuclear matrix protein, termed NM-200.4, was raised from the breast carcinoma cell line T-47D. Weidner et al., *Am. J. Path.* 138: 1293-98 (1991). This antibody reacts strongly with human breast carcinoma specimens as well as specimens from lung, thyroid, and ovarian cancers, but does not react with normal epithelial cells of similar origin.

WO 97/16206

-3-

PCT/US96/17119

raising the possibility of the use of certain anti-NMP antibodies as diagnostic tools.

In co-pending application Serial No. 08/015,624, the entire contents of which are incorporated by reference herein, it has been demonstrated with the Dunning rat model of prostate cancer that nuclear matrix protein composition is altered when comparing the normal dorsal prostate with the spontaneously arisen rat prostate adenocarcinomas. When human prostate samples were examined, nuclear matrix proteins were identified that (1) were present only in the normal prostate and were missing in both prostate cancer and benign prostatic hyperplasia (BPH) (normal pattern), (2) were found only in the prostate cancer cells and missing in the normal prostate and BPH (prostate cancer pattern), and (3) were found in both normal and BPH samples but were absent from prostate cancers.

No nuclear matrix proteins have been identified heretofore, however, that are linked specifically to bladder cancer.

SUMMARY OF THE INVENTION

The present invention relates to nuclear matrix proteins that are able to differentiate cancerous bladder cells from normal bladder cells, polynucleotide sequences encoding them, and their methods of use. Six proteins, respectively designated BLCA-1, BLCA-2, BLCA-3, BLCA-4, BLCA-5 and BLCA-6, have been discovered that are present in all cancerous bladder cells that are not present in the normal bladder cells, and three proteins (referred to as BLNL-1, BLNL-2, and BLNL-3) have been discovered that are unique to normal bladder tissue. These proteins are useful for diagnosing and producing treatments for cell proliferative disorders of the bladder.

WO 97/16206

- 4 -

PCT/US96/17119

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

According to one aspect, the present invention is directed to a purified nuclear matrix protein or a fragment thereof, which is present in normal bladder cells but absent in cancerous bladder cells, or which is absent in normal bladder cells but present in cancerous bladder cells. In particular, the present invention relates to a protein that is present in normal bladder cells but absent in cancerous bladder cells selected from the group consisting of BLNL-1, BLNL-2, and BLNL-3. In addition, the present invention relates to a protein that is absent in normal bladder cells but present in cancerous bladder cells selected from the group consisting of BLCA-1, BLCA-2, BLCA-3, BLCA-4, BLCA-5, and BLCA-6.

Another embodiment of the present invention is a purified polynucleotide sequence encoding the above identified NMPs or NMP fragments of the preceding embodiment. Another embodiment is a purified polynucleotide sequence which hybridizes to the polynucleotide sequence encoding the above-mentioned NMPs or NMP fragments.

Another embodiment is a host cell transformed with a polynucleotide sequence encoding the above-mentioned NMPs or NMP fragments. Transformation of a host cell with recombinant DNA may be carried out by conventional techniques known in the art. Where the host is prokaryotic, such as *E. coli*, competent cells which are capable of DNA uptake can be prepared from cells harvested after the exponential growth phase and subsequently treated by the CaCl_2 method by procedures well known in the art. Alternatively, MgCl_2 or RbCl can be used. Transformation can also be performed after forming a protoplast of the host cell or by electroporation.

When the host is a eukaryote, such methods of transfection of DNA as calcium phosphate co-precipitates,

WO 97/16206

-5-

PCT/US96/17119

conventional mechanical procedures such as micro-injection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors may be used. Eukaryotic cells can also be cotransformed with DNA sequences encoding the NMPs of the invention, and a second foreign DNA molecule encoding a selectable phenotype, such as the herpes simplex thymidine kinase gene. Another method is to use a eukaryotic viral vector, such as simian virus 40 (SV40) or bovine papilloma virus, to transiently infect or transform eukaryotic cells and express the protein. EUKARYOTIC VIRAL VECTORS Gluzman (ed.), Cold Spring Harbor Laboratory, 1982.

Isolation and purification of the NMPs or NMP fragments expressed by a transformed host may be carried out by conventional means including preparative chromatography and immunological separations involving monoclonal or polyclonal antibodies. Antibodies provided in the present invention are immunoreactive with NMP polypeptide or fragments thereof.

Another embodiment is a recombinant expression vector containing the above-mentioned polynucleotide sequences. Preferably, the vector is a virus. Preferred viruses are RNA viruses and preferred RNA viruses are retroviruses. Another preferred vector is a liposome, preferably a target-specific liposome which may be targeted with, for example, an antibody or ligand. Another preferred vector is a plasmid.

Another embodiment is an antibody which binds to the above-mentioned NMPs or NMP fragments. The antibody may be polyclonal or monoclonal.

Another embodiment is a method for detecting a cell proliferative disorder in a subject, comprising contacting a cellular component from the subject with an antibody or nucleic acid probe which binds to a cellular component associated with the cell proliferative disorder. Preferably, the cellular component is taken from the subject's bladder and is preferably nucleic acid. Preferably, the nucleic acid is DNA encoding the

WO 97/16206

-6-

PCT/US96/17119

above-mentioned NMPs or NMP fragments. Also preferred as a nucleic acid is RNA. Another preferred cellular component is the above-mentioned NMPs or NMP fragments.

Preferably, the nucleic acid probe specifically
5 hybridizes to the above-mentioned cellular component. When the reagent is a nucleic acid probe, it is preferably detectably labeled. Preferred labels include a radioisotope, a bioluminescent compound, a chemiluminescent compound, a fluorescent compound, a
10 metal chelate, and an enzyme.

Alternatively, if the cellular component is an NMP or NMP fragment, then an antibody is used which specifically binds to the NMP or NMP fragment. As noted above, the antibody may be monoclonal or polyclonal.

15 Another embodiment is a method of treating a cell proliferative disorder associated with a protein selected from the group consisting of BLCA-1, BLCA-2, BLCA-3, BLCA-4, BLCA-5, BLCA-6, BLNL-1, BLNL-2, and BLNL-3, comprising administering to a subject with the disorder
20 a therapeutically effective amount of an antisense polynucleotide sequence that blocks the sequences encoding the above-mentioned NMPs. In this embodiment, the treatment is designed to block the expression of one or more NMPs which give rise to the cell proliferative
25 disorder.

In an alternative method of treatment, instead of using an antisense polynucleotide sequence, a polynucleotide sequence is used which encodes one of the above-mentioned NMPs. In this embodiment, the treatment
30 is designed to provide the subject with one or more NMPs that prevent or ameliorate the cell proliferative disorder.

In another method of treatment, an antibody is administered to the subject which is capable of blocking
35 the function of one or more of the above NMPs.

Another embodiment is a method of gene therapy, comprising introducing into the cells of a host subject an expression vector comprising a polynucleotide sequence encoding one or more of the above-mentioned NMPs.

WO 97/16206

-7-

PCT/US96/17119

Preferably, the expression vector is introduced into the cells of the host subject *ex vivo*, yielding transformed cells, and the transformed cells then are reintroduced into the subject. A preferred expression vector for this purpose is an RNA virus, preferably a retrovirus.

Another embodiment of the present invention relates to a method for identifying a composition which blocks or enhances the function of a bladder cell NMP. The inventive method comprises:

- (a) incubating NMP-containing bladder cells with a test composition under conditions that allow the bladder cells and test composition to interact,
and then
- (b) measuring whether the test composition blocks or enhances the function of the bladder cell NMP.

Another embodiment is a kit for detecting a cell-proliferative disorder of the bladder comprising a nucleic acid probe that binds to a polynucleotide sequence encoding one of the above-mentioned NMPs. Preferably, the probe is labeled for ease of detection with a label as described above. Alternatively, the kit may comprise an antibody which specifically binds to one of the above-mentioned NMPs. Still another alternative is to use an oligonucleotide primer in the kit that permits amplification of a target polynucleotide sequence encoding one of the above-mentioned NMPs, for example, by polymerase chain reaction (PCR) amplification.

The NMPs of the invention include fragments and conservatively substituted variants thereof. Minor modifications of the NMP primary amino acid sequence may result in proteins which have substantially equivalent activity as compared to the NMP polypeptide described herein. Such modifications may be deliberate, as by site-directed mutagenesis, or may be spontaneous. Such modifications include deletion of non-essential amino acids. All of the polypeptides produced by these modifications are included herein as long as the

WO 97/16206

- 8 -

PCT/US96/17119

biological activity of the native NMP still exists. Further, deletion of one or more amino acids can also result in a modification of the structure of the resultant molecule without significantly altering its biological activity. This can lead to the development of a smaller active molecule which would have broader utility.

The term "conservative substitution" as used herein denotes the replacement of an amino acid residue by a structurally similar residue. Examples of conservative substitutions include the substitution of one hydrophobic residue such as isoleucine, valine, leucine or methionine for another, or the substitution of one polar residue for another, such as the substitution of arginine for lysine, glutamic for aspartic acids, or glutamine for asparagine, and the like.

Peptides of the invention can be synthesized by the well known solid phase peptide synthesis methods described, for example, by Merrifield, *J. Am. Chem. Soc.* 85: 2149 (1962), and by Stewart and Young, *SOLID PHASE PEPTIDES SYNTHESIS* 27-62 (Freeman Publ., 1969).

The polyclonal and monoclonal antibodies of the invention are immunoreactive with the NMPs or immunogenic fragments of the NMPs. If desired, polyclonal antibodies can be further purified, for example, by binding to and elution from a matrix to which NMP polypeptide is bound or by utilizing common nuclear matrix proteins to selectively remove non-specific antibodies. Antibody which consists essentially of pooled monoclonal antibodies with different epitopic specificities, as well as distinct monoclonal antibody preparations are provided. The term "antibody" as used in this invention includes intact molecules as well as fragments thereof, such as Fab and F(ab')₂ fragments, which are functionally capable of binding an epitopic determinant of an NMP.

A preferred method for the identification and isolation of antibody binding domains which exhibit binding with NMP is the bacteriophage λ vector system. This vector system has been used to express a combina-

WO 97/16206

-9-

PCT/US96/17119

torial library of Fab fragments from the mouse antibody repertoire in *Escherichia coli*, see Huse et al., Science 246: 1275-81 (1989), and from the human antibody repertoire. Mullinax et al., Proc. Nat'l Acad. Sci. USA 87: 8095-99 (1990).

As used herein, the term "cell-proliferative disorder" denotes malignant as well as non-malignant (or benign) disorders of the bladder. This term further encompasses hyperplastic disorders of the bladder. The cells comprising these proliferative disorders often appear morphologically and genotypically to differ from the surrounding normal tissue. As noted above, cell-proliferative disorders may be associated, for example, with expression or absence of expression of the NMPs of the invention. Expression of an NMP at an inappropriate time during the cell cycle or in an incorrect cell type may result in a cell-proliferative disorder. The NMP-encoding polynucleotide in the form of an antisense polynucleotide is useful in treating hyperplasia and malignancies of the bladder. When the cell-proliferative disorder is associated with NMP expression, (e.g., BLCA-1, 2, 3, 4, 5 and/or 6), an antisense NMP polynucleotide sequence or NMP binding antibody can be introduced into the bladder cells to block the expression and/or function of the NMP. Alternatively, when the cell proliferative disorder is associated with under-expression or expression of a mutant NMP polypeptide (e.g., BLNL 1-3), a polynucleotide sequence encoding the missing or under-expressed NMP can be introduced into the cell.

For purposes of the invention, an antibody or nucleic acid probe specific for an NMP may be used to detect the presence of the NMP polypeptide (in the case of an antibody probe) or polynucleotide (in the case of the nucleic acid probe) in biological fluids or tissues suspected of containing the NMP. Oligonucleotide primers based on any coding sequence region in the NMP sequence are useful for amplifying DNA or RNA, for example by PCR. Any specimen containing a detectable amount of antigen can be used. A preferred sample of this invention is

WO 97/16206

-10-

PCT/US96/17119

tissue taken from the bladder. Alternatively, biological fluids which may contain cells of the bladder may be used.

5 The term "subject" as used herein refers to mammals, preferably humans.

Another technique which may also result in greater sensitivity consists of coupling the probe to low molecular weight haptens. These haptens can then be specifically detected by means of a second reaction. For
10 example, it is common to use such haptens as biotin, which reacts with avidin, or dinitrophenol, pyridoxal, and fluorescein, which can react with specific antihapten antibodies.

The method for detecting a cell expressing a
15 particular NMP of the invention or a cell-proliferative disorder associated with an NMP, described above, can be utilized for detection of residual bladder cancer or other malignancies or benign hyperplasia conditions in a subject in a state of clinical remission. Additionally,
20 the method for detecting NMP polypeptide in cells is useful for detecting a cell-proliferative disorder by identifying cells expressing specific NMPs in comparison with NMPs expressed in normal cells. Using the method of the invention, NMP expression can be identified in a cell
25 and the appropriate course of treatment can be employed (e.g., NMP-encoding or antisense gene therapy, as well as conventional chemotherapy). Since the expression pattern of the NMPs of the invention vary with the stage of malignancy of a cell, a sample of bladder tissue can be
30 screened with a panel of NMP-specific reagents (e.g., nucleic acid probes or antibodies to NMPs) to detect NMP expression and diagnose the stage of malignancy of the cell.

The monoclonal antibodies of the invention are
35 suitable for use in immunoassays in which they can be utilized in liquid phase or bound to a solid phase carrier. In addition, the monoclonal antibodies in these immunoassays can be detectably labeled in various ways. Examples of types of immunoassays which can utilize

WO 97/16206

-11-

PCT/US96/17119

monoclonal antibodies of the invention are competitive and non-competitive immunoassays in either a direct or indirect format. Examples of such immunoassays are the radioimmunoassay (RIA) and the sandwich (immunometric) assay. Detection of the antigens using the monoclonal antibodies of the invention can be performed utilizing immunoassays which are run in either the forward, reverse, or simultaneous modes, including immunohistochemical assays on physiological samples. Alternatively, the antibody of the invention can be used to detect NMPs present in electrophoretically dispersed gel protocols such as Western blots and two-dimensional gels.

The monoclonal antibodies of the invention can be bound to many different carriers and used to detect the presence of NMP. Examples of well-known carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, agaroses and magnetite. The nature of the carrier can be either soluble or insoluble for purposes of the invention.

In performing the assays it may be desirable to include certain "blockers" in the incubation medium (usually added with the labeled soluble antibody). The "blockers" are added to assure that non-specific proteins, proteases, or anti-heterophilic immunoglobulins to anti-NMP immunoglobulins present in the experimental sample do not cross-link or destroy the antibodies on the solid phase support, or the radiolabeled indicator antibody, to yield false positive or false negative results. The selection of "blockers" therefore may add substantially to the specificity of the assays described in the present invention.

It has been found that a number of nonrelevant (i.e., nonspecific) antibodies of the same class or subclass (isotype) as those used in the assays (e.g., IgG1, IgG2a, IgM, etc.) can be used as "blockers." The "blockers" are used at a level high enough to maintain the proper sensitivity yet inhibit any unwanted

WO 97/16206

-12-

PCT/US96/17119

interference by mutually occurring cross reactive proteins in the specimen (normally 1-100 $\mu\text{g}/\mu\text{l}$).

In this description, the term "epitope" denotes any determinant capable of specific interaction with the monoclonal antibodies of the invention. Epitopic determinants usually comprise chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics.

In using the monoclonal antibodies of the invention for the *in vivo* detection of antigen, the detectably labeled monoclonal antibody is given in a dose which is diagnostically effective. The term "diagnostically effective" means that the amount of detectably labeled monoclonal antibody is administered in sufficient quantity to enable detection of the site having the NMP antigen for which the monoclonal antibody is specific. The dosage of detectably labeled monoclonal antibody for *in vivo* diagnosis will vary depending on such factors as age, sex, and extent of disease of the individual. The dosage of monoclonal antibody can vary from about 0.001 mg/m^2 , to about 500 mg/m^2 , preferably 0.1 mg/m^2 to about 200 mg/m^2 , most preferably about 0.1 mg/m^2 to about 10 mg/m^2 . Such dosages may vary, for example, depending on whether multiple injections are given, tumor burden, and other factors.

For *in vivo* diagnostic imaging, the type of detection instrument available is a major factor in selecting a given radioisotope. The radioisotope chosen must have a type of decay which is detectable for a given type of instrument. Still another important factor in selecting a radioisotope for *in vivo* diagnosis is that the half-life of the radioisotope be long enough so that it is still detectable at the time of maximum uptake by the target, but short enough so that deleterious radiation with respect to the host is minimized. Ideally, a radioisotope used for *in vivo* imaging will lack a particle emission, but produce a large number of

WO 97/16206

-13-

PCT/US96/17119

photons in the 140-250 keV range, which may be readily detected by conventional gamma cameras.

For in vivo diagnosis, radioisotopes may be bound to immunoglobulin either directly or indirectly by using an intermediate functional group. Intermediate functional groups which often are used to bind radioisotopes which exist as metallic ions to immunoglobulins are the bifunctional chelating agents such as diethylenetriamine-pentacetic acid (DTPA) and ethylenediaminetetraacetic acid (EDTA) and similar molecules. Typical examples of metallic ions which can be bound to the monoclonal antibodies of the invention are ^{111}In , ^{97}Ru , ^{67}Ga , ^{68}Ga , ^{72}As , ^{89}Zr , and ^{201}Tl .

The monoclonal antibodies of the invention can also be labeled with a paramagnetic isotope for purposes of in vivo diagnosis, as in magnetic resonance imaging (MRI) or electron spin resonance (ESR). In general, any conventional method for visualizing diagnostic imaging can be utilized. Usually gamma and positron emitting radioisotopes are used for camera imaging and paramagnetic isotopes for MRI. Elements which are particularly useful in such techniques include ^{157}Gd , ^{55}Mn , ^{162}Dy , ^{52}Cr , and ^{56}Fe .

The monoclonal antibodies of the invention can be used to monitor the course of amelioration of an NMP-associated cell-proliferative disorder. Thus, by measuring the increase or decrease in the number of cells expressing a NMP or changes in NMP present in various body fluids, such as ejaculate or urine, it would be possible to determine whether a particular therapeutic regimen aimed at ameliorating the disorder is effective.

The monoclonal antibodies of the invention can also be used, alone or in combination with effector cells, see Douillard et al., *Hybridoma* 5 (Supp. 1): S139 (1986), for immunotherapy in an animal having a cell proliferative disorder which expresses NMP polypeptide with epitopes reactive with the monoclonal antibodies of the invention.

When used for immunotherapy, the monoclonal antibodies of the invention may be unlabeled or attached to

WO 97/16206

-14-

PCT/US96/17119

a therapeutic agent. These agents can be coupled either directly or indirectly to the monoclonal antibodies of the invention. One example of indirect coupling is by use of a spacer moiety. These spacer moieties, in turn, can be either insoluble or soluble, see Diener et al., *Science* 231: 148 (1986), and can be selected to enable drug release from the monoclonal antibody molecule at the target site. Examples of therapeutic agents which can be coupled to the monoclonal antibodies of the invention for immunotherapy are drugs, radioisotopes, lectins, and toxins.

The drugs which can be conjugated to the monoclonal antibodies of the invention include non-proteinaceous as well as proteinaceous drugs. The terms "non-proteinaceous drugs" encompasses compounds which are classically referred to as drugs, for example, mitomycin C, daunorubicin, vinblastine, and others used to treat cancer.

The proteinaceous drugs with which the monoclonal antibodies of the invention can be joined include immunomodulators and other biological response modifiers. The term "biological response modifiers" encompasses substances which are involved in modifying the immune response in such manner as to enhance the destruction of an NMP-associated tumor for which the monoclonal antibodies of the invention are specific. Examples of immune response modifiers include such compounds as lymphokines. Lymphokines include tumor necrosis factor, the interleukins, lymphotoxin, macrophage activating factor, migration inhibition factor, colony stimulating factor, and interferon. Interferons with which the monoclonal antibodies of the invention can be labeled include alpha-interferon, beta-interferon and gamma-interferon and their subtypes.

In using radioisotopically conjugated monoclonal antibodies of the invention for immunotherapy certain isotopes may be more preferable than others depending on such factors as tumor cell distribution as well as isotope stability and emission. If desired, the tumor

WO 97/16206

-15-

PCT/US96/17119

cell distribution can be evaluated by the *in vivo* diagnostic techniques described above. Depending on the cell proliferative disease some emitters may be preferable to others. In general, alpha and beta particle-emitting radioisotopes are preferred in immunotherapy. For example, if an animal has solid tumor foci a high energy beta emitter capable of penetrating several millimeters of tissue, such as ^{90}Y , may be preferable. On the other hand, if the cell proliferative disorder consists of simple target cells, as in the case of leukemia, a short range, high energy alpha emitter, such as ^{212}Bi , may be preferable. Examples of radioisotopes which can be bound to the monoclonal antibodies of the invention for therapeutic purposes are ^{125}I , ^{131}I , ^{90}Y , ^{67}Cu , ^{212}Bi , ^{211}At , ^{212}Pb , ^{47}Sc , ^{109}Pd , ^{65}Zn , and ^{188}Re .

Lectins are proteins, usually isolated from plant material, which bind to specific sugar moieties. Many lectins are also able to agglutinate cells and stimulate lymphocytes. Ricin is a toxic lectin which has been used immunotherapeutically. The alpha-peptide chain of ricin, which is responsible for toxicity, may be bound to the antibody of the invention to enable site specific delivery of the toxic effect.

Toxins are poisonous substances produced by plants, animals, or microorganisms, that, in sufficient dose, are often lethal. Diphtheria toxin is a substance produced by *Corynebacterium diphtheria* which can be used therapeutically. This toxin consists of an alpha and beta subunit which under proper conditions can be separated. The toxic A component can be bound to an antibody and used for site specific delivery to a NMP bearing cell.

The monoclonal antibodies of the invention can be used in combination with alpha-interferon. This treatment modality enhances monoclonal antibody targeting of carcinomas by increasing the expression of monoclonal antibody reactive antigen by the carcinoma cells. Greiner et al., *Science* 235: 895 (1987). Alternatively, the monoclonal antibody of the invention can be used, for

WO 97/16206

-16-

PCT/US96/17119

example, in combination with gamma-interferon to thereby activate and increase the expression of Fc receptors by effector cells which, in turn, results in an enhanced binding of the monoclonal antibody to the effector cell and killing of target tumor cells.

It is also possible to utilize liposomes with the monoclonal antibodies of the invention in their membrane to specifically deliver the liposome to the tumor expressing NMP. These liposomes can be produced such that they contain, in addition to the monoclonal antibody, such immunotherapeutic agents as those described above which would then be released at the tumor site. Wolff et al., *Biochemical et Biophysical Acta* 802: 259 (1984).

The dosage ranges for the administration of monoclonal antibodies of the invention are those large enough to produce the desired effect in which the symptoms of the malignant disease are ameliorated. The dosage should not be so large as to cause adverse side effects, such as unwanted cross-reactions, anaphylactic reactions, and the like. Generally, the dosage will vary with the age, condition, sex and extent of the disease in the patient and can be determined by one of skill in the art. The dosage can be adjusted by the individual physician in the event of any complication. Dosage can vary from about 0.1 mg/kg to about 2000 mg/kg, preferably about 0.1 mg/kg to about 500 mg/kg, in one or more dose administrations daily, for one or several days. Generally, when the monoclonal antibodies of the invention are administered conjugated with therapeutic agents, lower dosages, comparable to those used for in vivo diagnostic imaging, can be used.

The monoclonal antibodies of the invention can be administered parenterally by injection or by gradual perfusion over time. The monoclonal antibodies of the invention can be administered intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, or transdermally, alone or in combination with effector cells.

WO 97/16206

-17-

PCT/US96/17119

The present invention also provides a method for treating a subject with an NMP-associated cell-proliferative disorder using an NMP nucleotide sequence. An NMP nucleotide sequence which may encode a suppressor polypeptide may be under-expressed as compared to expression in a normal cell, therefore it is possible to design appropriate therapeutic or diagnostic techniques directed to this sequence. Thus, where a cell-proliferative disorder is associated with the expression of an NMP associated with malignancy, nucleic acid sequences that interfere with NMP expression at the translational level can be used. This approach utilizes, for example, antisense nucleic acid and ribozymes to block translation of a specific NMP mRNA, either by masking that mRNA with an antisense nucleic acid or by cleaving it with a ribozyme. In cases when a cell proliferative disorder or abnormal cell phenotype is associated with the under expression of NMP suppressor for example, nucleic acid sequences encoding NMP (sense) could be administered to the subject with the disorder.

Antisense nucleic acids are DNA or RNA molecules that are complementary to at least a portion of a specific mRNA molecule. Weintaub, *Scientific American*, 262: 40 (1990). In the cell, the antisense nucleic acids hybridize to the corresponding mRNA, forming a double-stranded molecule. The antisense nucleic acids interfere with the translation of the mRNA since the cell will not translate a mRNA that is double-stranded. Antisense oligomers of about 15 nucleotides are preferred, since they are easily synthesized and are less likely to be expressed than larger molecules when introduced into the target NMP-producing cell.

Ribozymes are RNA molecules possessing the ability to specifically cleave other single-stranded RNA in a manner analogous to DNA restriction endonucleases. Through the modification of nucleotide sequences which encode these RNAs, it is possible to engineer molecules that recognize specific nucleotide sequences in an RNA molecule and cleave it. Cech, *J. Amer. Med. Assn.* 260:

WO 97/16206

-18-

PCT/US96/17119

3030 (1988). A major advantage of this approach is that, because they are sequence-specific, only mRNAs with particular sequences are inactivated.

There are two basic types of ribozymes namely, 5 tetrahymena-type (Hasselhoff, *Nature*, 334:585, 1988) and "hammerhead"-type. Tetrahymena-type ribozymes recognize sequences which are four bases in length, while "hammerhead"-type ribozymes recognize base sequences 11-18 bases in length. The longer the recognition sequence, 10 the greater the likelihood that that sequence will occur exclusively in the target mRNA species. Consequently, hammerhead-type ribozymes are preferable to tetrahymena-type ribozymes for inactivating a specific mRNA species and 18-based recognition sequences are preferable to 15 shorter recognition sequences.

The present invention also provides gene therapy for the treatment of cell proliferative disorders which are mediated by NMP. Such therapy requires introduction of the appropriate NMP polynucleotide sequence (antisense or 20 encoding strand) into cells of subjects having the proliferative disorder. Delivery of antisense NMP polynucleotides can be achieved using a recombinant expression vector such as a chimeric virus or a liposome. Disorders associated with under-expression of an NMP or 25 expression of a cancer-associated NMP can be treated using gene therapy with the encoding or antisense nucleotide sequences, respectively.

Various viral vectors which can be utilized for gene therapy as taught herein include adenovirus, herpes 30 virus, vaccinia, or, preferably, an RNA virus such as a retrovirus. Preferably, the retroviral vector is a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus 35 (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a

WO 97/16206

-19-

PCT/US96/17119

selectable marker so that transduced cells can be identified and generated. By inserting an NMP sequence of interest into the viral vector along with another gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is rendered target specific. Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a sugar, a glycolipid, or a protein. Preferred targeting is accomplished by using an antibody to target the retroviral vector.

Since recombinant retroviruses are defective in one or more genes, they require assistance in order to produce infectious vector particles. Helper cell lines which have deletions of the packaging signal include but are not limited to Ψ 2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced.

Alternatively, NIH 3T3 or other tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes *gag*, *pol* and *env* by conventional calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The resulting cells release the retroviral vector into the culture medium.

Other targeted delivery systems for NMP antisense polynucleotides include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. Liposomes are artificial membrane vesicles which are useful as delivery vehicles in vitro and in vivo. It has been shown that large unilamellar vesicles (ULV), which range in size from 0.2-4.0 μ m can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA and intact virions can be encapsulated within the aqueous

WO 97/16206

-20-

PCT/US96/17119

interior and be delivered to cells in a biologically active form. Fraley et al., *Trends Biochem. Sci.* 6: 77 (1981).

5 The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic
10 strength, and the presence of divalent cations.

Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidyl-glycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebrosides,
15 and gangliosides. Particularly useful are diacyl-phosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine.
20

The targeting of liposomes has been classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific and organelle-specific.
25 Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries.
30 Active targeting, on the other hand, involves alteration of the liposome by coupling the liposome to a specific ligand such as a monoclonal antibody, sugar, glycolipid, or protein, or by changing the composition or size of the liposome in order to achieve targeting to organs and cell
35 types other than the naturally occurring sites of localization.

The surface of the targeted delivery system may be modified in a variety of ways. In the case of a hyposomal targeted delivery system, lipid groups can be

WO 97/16206

-21-

PCT/US96/17119

incorporated into the lipid bilayer of the liposome in order to maintain the targeting ligand in stable association with the liposomal bilayer. Various linking groups can be used for joining the lipid chains to the
5 targeting ligand.

In general, the compounds bound to the surface of the targeted delivery system will be ligands and receptors which will allow the targeted delivery system to find and "home in" on the desired cells. A ligand may
10 be any compound of interest which will bind to another compound, such as a receptor.

In general, surface membrane proteins which bind to specific effector molecules are referred to as receptors. In the present invention, antibodies of the invention are preferred receptors. Antibodies can be used to target
15 liposomes to specific cell-surface ligands, in this case the NMPs of choice. Preferably, the target tissue is bladder tissue. A number of procedures can be used to covalently attach either polyclonal or monoclonal
20 antibodies to a liposome bilayer. Antibody-targeted liposomes can include monoclonal or polygonal antibodies or fragments thereof such as Fab, or F(ab')₂, as long as they bind efficiently to an the antigenic epitope on the target cells.

Preparations for parenteral administration include sterile aqueous or nonaqueous solutions, suspensions, and emulsions. Examples of non aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as
30 ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's
35 intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such an those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for

WO 97/16206

-22-

PCT/US96/17119

example, antimicrobials, anti-oxidants, chelating agents and inert gases and the like.

The invention also relates to a method for preparing a medicament or pharmaceutical composition comprising the polynucleotides or the monoclonal antibodies of the invention, the medicament being used for therapy of NMP associated cell proliferative disorders.

The invention is further illustrated by, though in no way limited to, the following examples.

10 Tissue Selection

Normal and tumor bladder tissue samples were obtained from patients undergoing surgery for bladder cancer at the University of Pittsburgh Medical Center. Normal bladders were obtained from the Center for Organ Recovery and Education (CORE) via Dr. Michael J. Becich, Department of Pathology, University of Pittsburgh. These bladders provide the opportunity to study nuclear matrix composition and staining in the normal organ which were compared with alterations that occur during transformation from normal to tumor. Samples were only utilized that could clearly be identified by the pathologist as containing approximately pure populations of the stated cell type.

Nuclear Matrix Preparation

25 The nuclear matrix proteins were isolated from the bladder tissue and tumors selected above as taught in Fey, et al., *J. Cell Biol.*, 98:1973-1984, 1988 and Getzenberg, et al., *Cancer Res.*, 51:6514-6520, 1991. The tissue pieces were minced into small (1 mm³) pieces and homogenized with a Teflon pestle on ice with 0.5% Triton X-100 in a solution containing 2 mM vanadyl ribonucleoside (RNase inhibitor) to release the lipids and soluble proteins. Extracts were then filtered through a 350 micron nylon mesh and extracted with 0.25 M ammonium sulfate to release the soluble cytoskeletal elements. Dnase treatment at 25°C was used to remove the soluble chromatin. The remaining fractions contained

WO 97/16206

PCT/US96/17119

-23-

intermediate filaments and nuclear matrix proteins. This fraction was then disassembled with 8 M urea, and the insoluble components, which consisted principally of carbohydrates and extracellular matrix components, were pelleted. The urea was dialyzed out, and the intermediate filaments were allowed to reassemble and removed by centrifugation. The nuclear matrix proteins were then ethanol precipitated. All solutions contained freshly prepared 1 mM phenylmethylsulfonylfluoride (PMSF) to inhibit serine proteases, 0.3 μ M aprotonin, 1 μ M leupeptin and 1 μ M pepstatin. Antibodies to proteins of this fraction were prepared and demonstrated to be localized exclusively in the nucleus and isolated nuclear matrix fraction. The protein composition was determined by resuspending the proteins in 0.1 N sodium hydroxide and utilizing the Coomassie Plus protein assay reagent kit (Pierce, Rockford, IL) with bovine serum albumin (BSA) as a standard.

For two-dimensional gel electrophoresis, the ethanol precipitated NMPs were dissolved in a sample buffer consisting of 9 M urea, 65 mM 3-[(3-Cholamidopropyl)-dimethyl-ammonio]-1-propanesulfonate (CHAPS), 2.2% ampholytes and 140 mM dithiothreitol. The final pellet containing NMPs represented less than 1% of the total cellular proteins.

High Resolution Two-Dimensional Electrophoresis

High resolution two-dimensional gel electrophoresis was carried out utilizing the Investigator 2-D gel system (Oxford Glycosystems, Bedford, MA). Briefly, one-dimensional isoelectric focusing was carried out for 18,000 V-h using 1-mm x 18-cm tube gels after 1.5 h of prefocusing. The tube gels were extruded and placed on top of 1-mm sodium dodecyl sulfate Duracryl (Oxford Glycosystems, Bedford, MA) high tensile strength polyacrylamide electrophoresis slab gels, and the gels were electrophoresed with 12°C constant temperature regulation for approximately 5 hours. Gels were fixed with 50% methanol and 10% acetic acid. After thorough

WO 97/16206

-24-

PCT/US96/17119

rinsing and rehydration, gels were treated with 5% glutaraldehyde and 5 mM dithiothreitol after buffering with 50 mM phosphate (pH 7.2). The gels were stained with silver stain (Accurate Chemical Co., Inc., Westbury, NY) or transferred to PVDF (Immobilon, Millipore Corporation) as follows. Fifty micrograms of nuclear matrix protein were loaded for each gel. Protein molecular weight standards were Silver Standards from Diversified Biotechnology (Newton Centre, MA).

10 Isoelectric points were determined using carbamylated standards from Gallaro-Schlesiwger, Inc. (Carle Place, NY) and Sigma Chemical Co. (St. Louis, MO). Multiple gels were run for each sample, and multiple samples run at different times. Only protein spots clearly and

15 reproducibly observed in all the gels of a sample type were counted as actually representing the nuclear matrix components. The gels were analyzed using the BioImage Electrophoresis Analysis System (BioImage, Ann Arbor, MI) which matches protein spots between gels and databases

20 the gels and protein spots.

WO 97/16206

-25-

PCT/US96/17119

The following NMPs were identified:

Table I

Proteins Associated With Human Bladder Cancer

		<u>Molecular Weight (kD)</u>	<u>pI</u>
5	BLCA-1	72	7.70
	BLCA-2	40	7.50
	BLCA-3	39	6.27
	BLCA-4	37	6.24
	BLCA-5	29	5.80
10	BLCA-6	22	8.00

Proteins Associated with Normal Human Bladder

		<u>Molecular Weight (kD)</u>	<u>pI</u>
	BLNL-1	70	6.09
	BLNL-2	66	5.84
15	BLNL-3	66	5.80

The designation of each protein above is shown in the gels displayed in Figures 1-5. In addition, the following preliminary sequence data has been obtained: BLCA-1 includes the amino acid sequence LAKIVL. BLCA-4 includes the amino acid sequences EISQLNAG and VYEDIMQK. BLCA-6 includes the amino acid sequence SLDLDLIIAEVK.

While the invention has been described in detail and with reference to specific embodiments thereof, it will be apparent to one skilled in the art that various changes and modifications can be made therein without departing from the spirit and scope of the invention.

WO 97/16206

-26-

PCT/US96/17119

We claim:

1. A purified nuclear matrix protein or a fragment thereof, which is present in normal bladder cells but absent in cancerous bladder cells, or which is absent in
5 normal bladder cells but present in cancerous bladder cells.
2. The protein or fragment as claimed in claim 1, wherein the protein is selected from the group consisting of BLNL-1, BLNL-2, and BLNL-3.
- 10 3. The protein or fragment of claim 1, wherein the protein is selected from the group consisting of BLCA-1, BLCA-2, BLCA-3, BLCA-4, BLCA-5, and BLCA-6.
4. A purified polynucleotide sequence encoding a protein or fragment of claim 1.
- 15 5. A purified polynucleotide sequence which hybridizes to the polynucleotide sequence of claim 4.
6. A host cell transformed with the polynucleotide of claim 4.
7. A recombinant expression vector containing the
20 polynucleotide of claim 4.
8. The vector of claim 7, wherein the vector is a virus.
9. The vector of claim 8, wherein the virus is an RNA virus.
- 25 10. The vector of claim 9, wherein the RNA virus is a retrovirus.
11. The vector of claim 7, wherein the vector is a liposome.

WO 97/16206

PCT/US96/17119

-27-

12. The vector of claim 11, wherein the liposome is target- specific.
13. The vector of claim 12, wherein the liposome is targeted with an antibody.
- 5 14. The vector of claim 7, wherein the vector is a plasmid.
15. An antibody which binds to the protein of claim 1.
- 10 16. A method for detecting a cell proliferative disorder in a subject, comprising contacting a cellular component from the subject with a reagent which binds to a cellular component associated with a cell proliferative disorder.
- 15 17. The method of claim 16, wherein the cellular component is taken from the subject's bladder.
18. The method of claim 17, wherein the cellular component is nucleic acid.
19. The method of claim 18, wherein the nucleic acid is DNA.
- 20 20. The method of claim 18, wherein the nucleic acid is RNA.
21. The method of claim 17, wherein the cellular component is protein.
22. The method of claim 16, wherein the reagent is a probe.
- 25 23. The method of claim 22, wherein the probe is nucleic acid.

WO 97/16206

-28-

PCT/US96/17119

24. The method of claim 22, wherein the probe is an antibody.

25. The method of claim 24, wherein the antibody is polyclonal.

5 26. The method of claim 24, wherein the antibody is monoclonal.

27. The method of claim 22, wherein the probe is detectably labeled.

10 28. The method of claim 27, wherein the label is selected from the group consisting of a radioisotope, a bioluminescent compound, a chemiluminescent compound, a fluorescent compound, a metal chelate, and an enzyme.

15 29. A method of treating a cell proliferative disorder associated with a protein selected from the group consisting of BLCA-1, BLCA-2, BLCA-3, BLCA-4, BLCA-5, BLCA-6, BLNL-1, BLNL-2, and BLNL-3, comprising administering to a subject with the disorder a therapeutically effective amount of reagent which blocks or enhances the function of the protein.

20 30. The method of claim 29, wherein the reagent is an antisense polynucleotide sequence.

25 31. The method of claim 29, wherein the reagent is a polynucleotide sequence encoding a protein selected from the group consisting of BLCA-1, BLCA-2, BLCA-3, BLCA-4, BLCA-5, BLCA-6, BLNL-1, BLNL-2, and BLNL-3.

32. The method of claim 29, wherein the reagent is an antibody.

33. The method of claim 32, wherein the antibody is monoclonal.

WO 97/16206

-29-

PCT/US96/17119

34. The method of claim 29, wherein the cell proliferative disorder is in bladder tissue.

35. A method of gene therapy, comprising introducing into the cells of a host subject an expression vector comprising a nucleotide sequence encoding a protein of claim 1.

36. The method of claim 35, wherein the expression vector is introduced into the cells of the host subject *ex vivo*, yielding transformed cells, and the transformed cells then are reintroduced into the subject.

37. The method of claim 36, wherein the expression vector is an RNA virus.

38. The method of claim 37, wherein the RNA virus is a retrovirus.

39. The method of claim 36, wherein the subject is human.

40. A method for identifying a composition which blocks or enhances the function of an NMP of bladder cells, which method comprises:

(a) incubating NMP-containing bladder cells with a test composition under conditions that allow the bladder cells and test composition to interact,

and then

(b) measuring whether the test composition causes blocking or enhancement of the function of an NMP of the bladder cells.

41. The method of claim 40, wherein the NMP is selected from the group consisting of BLCA-1, BLCA-2, BLCA-3, BLCA-4, BLCA-5, BLCA-6, BLNL-1, BLNL-2, and BLNL-3.

WO 97/16206

-30-

PCT/US96/17119

42. The method of claim 40, wherein the effect being measured is blocking of the function of the NMP.

43. The method of claim 40, wherein the effect being measured is enhancement of the function of the NMP.

5 44. A kit useful for the detection of a cell-proliferative disorder associated with a protein of claim 1, said kit comprising a probe for identifying a protein or fragment of claim 1 or a polynucleotide sequence encoding a protein or fragment of claim 1.

10 45. The kit of claim 44, wherein the probe is an antibody capable of binding to the protein or fragment.

46. The kit of claim 44, wherein the probe is a polynucleotide probe capable of hybridizing with the polynucleotide sequence encoding the protein or fragment.

15 47. The kit of claim 46, further comprising a container containing an oligonucleotide primer for amplification of the polynucleotide sequence encoding the protein or fragment.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/17119

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 38/28; C07K 1/00; C07H 21/02, 21/04; C12N 5/00, 15/00; G01N 33/53
US CL : 435/7.1, 240.2, 320.1, 810; 514/4.4; 530/350; 536/23.1, 24.31
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/7.1, 240.2, 320.1, 810; 514/4.4; 530/350; 536/23.1, 24.31

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MEDLINE, BIOSIS, EMBASE, CAPLUS, WPIDS
nuclear matrix protein/bladder/prostate/cancer

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4,882,268 A (PENMAN ET AL.) 21 November 1989, see entire document.	1-47
Y	GETZENBERG et al. Identification of nuclear matrix proteins in the cancer and normal rat prostate. Cancer Research. 15 December 1991, Vol. 51, pages 6514-6520, see entire document.	1-47
Y	CUPO, J.F. Electrophoretic analysis of nuclear matrix proteins and the potential clinical applications. Journal of Chromatography. 1991, Vol. 569, pages 389-406, see entire document.	1-3, 15-28, 40-47

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 DECEMBER 1996

Date of mailing of the international search report

17 JAN 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RAY F. EBERT

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORTInternational application No.
PCT/US96/17119**C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---	GORDON et al. Altered extracellular matrices influence cellular processes and nuclear matrix organizations of overlying human bladder urothelial cells. Cancer Research. 15 October 1993, Vol. 53, pages 4971-4977, see entire document.	1-3, 15-17, 21, 22, 24-28
Y		----- 29-34, 40-47
Y	WO 93/09437 A2 (MATRITECH, INC.) 13 May 1993, see entire document.	1-47
Y	WO 95/16919 A2 (MATRITECH, INC.) 22 June 1995, see entire document.	1-47
Y	WO 87/03910 A1 (MASSACHUSETTS INSTITUTE OF TECHNOLOGY) 02 July 1987, see entire document.	1-47
Y	WO 94/00573 A1 (MATRITECH, INC.) 06 January 1994, see entire document.	1-47
Y	WO 94/18222 A1 (THE JOHNS HOPKINS UNIVERSITY SCHOOL OF MEDICINE) 18 August 1994, see entire document.	1-47